



## Metal Impurity transport control in JET H-mode plasmas with central Ion Cyclotron Radiofrequency Heating

**Valisa, M.; Carraro, L.; Predebon, I.; Puiatti, M.E.; Angioni, C.; Coffey, I.; Giroud, C.; Taroni, L. Lauro; Alper, B.; Baruzzo, M.**

*Total number of authors:*  
19

*Publication date:*  
2010

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

### *Citation (APA):*

Valisa, M. (Author), Carraro, L. (Author), Predebon, I. (Author), Puiatti, M. E. (Author), Angioni, C. (Author), Coffey, I. (Author), Giroud, C. (Author), Taroni, L. L. (Author), Alper, B. (Author), Baruzzo, M. (Author), Belo daSilva, P. (Author), Buratti, P. (Author), Garzotti, L. (Author), Van Eester, D. (Author), Lerche, E. (Author), Mantica, P. (Author), Naulin, V. (Author), Tala, T. (Author), & Tsalias, M. (Author). (2010). Metal Impurity transport control in JET H-mode plasmas with central Ion Cyclotron Radiofrequency Heating. Sound/Visual production (digital)

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Metal Impurity Transport Control in JET H-mode Plasmas with Central *ICRH*

M. Valisa, M.E. Puiatti

L. Carraro<sup>1</sup>, I. Predebon<sup>1</sup>, C. Angioni<sup>2</sup>, I. Coffey<sup>3</sup>, C. Giroud<sup>4</sup>, L. Lauro Taroni<sup>1</sup>, B. Alper<sup>4</sup>, M. Baruzzo<sup>1</sup>, P. Belo da Silva<sup>5</sup>, P. Buratti<sup>6</sup>, L. Garzotti<sup>4</sup>, D. Van Eester<sup>7</sup>, E. Lerche<sup>7</sup>, P. Mantica<sup>8</sup>, V. Naulin<sup>9</sup>, T. Tala<sup>10</sup>, M. Tsalas<sup>11</sup> and JET-EFDA contributors<sup>12</sup>

*1* Consorzio RFX – Associazione EURATOM-ENEA sulla Fusione, Padova, Italy

*2* Max Planck Institut für Plasmaphysik, EURATOM-IPP Association, D-85748 Garching, Germany

*3* Department of Physics, Queen's University, Belfast, United Kingdom

*4* EURATOM/UKAEA Fusion Association, Culham Science Centre, Abingdon, OX14 3DB, UK

*5* EURATOM/IST 5Fusion Association, Instituto de Plasmas e Fusão Nuclear, Av. Rovisco Pais 1049-001 Lisbon Portugal *6* EURATOM-ENEA Association, C.R. Frascati, CP 65, 00044 Frascati, Italy

*7* Association EURATOM-Belgian State, LPP-ERM/KMS, Partner in TEC, B-1000 Brussels, Belgium

*8* Associazione EURATOM-ENEA sulla Fusione, Via Cozzi, Milano, Italy

*9* Association EURATOM - Risoe DTU, Frederiksborgvej 399, 4000 Roskilde, Denmark

*10* Association Euratom-Tekes, VTT, P.O. Box 1000 FI-02044 VTT –Finland

*11* EFDA CSU Culham, Culham Science Centre, Abingdon, OX14 3DB, UK

*12* See Appendix of F. Romanelli et al., Proc. 22nd IAEA Fusion Energy Conf. 2008, Geneva, Switzerland

- We need to fully understand the behaviour of impurities in reactor relevant plasmas as in ITER and Demo plasma dilution in the core will be a figure of merit ( including fusion ashes).
- Also, as for many other parameters, impurities will need an active control system to guarantee stationarity of plasma conditions

- Source mechanisms and transport across SOL and pedestal important but beyond the scope of this work, **focused on core aspects**.
- For impurities in the core what really matters is the relationship between  $D_{\text{impurities}}$ ,  $D_{\text{fuel}}$  and  $\chi_{e,i}$ , since the relevant parameter is dilution.

Here we concentrate on  $D_{\text{impurities}}$

To measure impurity transport one powerful means is to **create impurity density perturbations as with laser ablation**. Modelling of the transient evolutions of the appropriate signals provides an estimate of the transport coefficients.

Model based on 1D continuity equation with  $\Gamma = -D\nabla n + vn$

+ accurate atomic physics to describe all of the ionization stages.

$$-\frac{\nabla n_z}{n_z} = -\frac{v}{D}$$

↓

Peaking factor

$>0$   
peaked profiles

$<0$   
hollow profiles

## • Turbulence and impurity radial transport

- *Curvature pinch (Perp Dynamics)* : inward  ~~$f(Z)$~~
- *Parallel Compressibility* : Outward for TEM  
Inward for ITG  $\sim Z/A$
- *Thermodiffusion* : Inward for TEM  $\sim 1/Z \rightarrow$  lower for high Z impurities  
Outward for ITG
- Curvature pinch changes sign with **magnetic shear** (Futatani)
- **Rotation and shear rotation** inward for TEM and Outward for ITG (Camenen)
- **Centrifugal and Coriolis forces** : outward advection (Clements/ Romanelli )
- **Electromagnetic effects**  $\sim 10\%$  of electrostatic ones (Hein)
- **RF induced ponderomotive forces** - affect similarly D and v ( Nordman)
- **High Impurity concentration**: significant in case of TEM (Fulop/ Moradi)

## • Neoclassical radial transport

$$\bullet \ v_{neo} / D_{neo} = Z_i \bullet (1/n \bullet dn/dr - H \bullet 1/T \bullet dT/dr) ; H \text{ approx } 1$$


RF ( electron heating) pump out effect observed in several other experiments:

C-mod, DIII-D, JT-60, TCV

and also in non axisymmetric devices (W7-AS)

...but only few example of successful explanation of radial flow reversal from  
the theory - AUG : TEM (Angioni et al. PPCF 2007)

JET discharges 58143 and 58142



*M.E. Puiatti et al  
PoP 13 2006*

*Steady state Ni profiles, calculated from exp  $v$  and  $D$*

Similar results obtained also in high density JET discharges

*M.E. Puiatti et al .Plas. Phys.Contr. Fus. 44(2002)1863*


A series of dedicated H mode discharges with a RF power scan + LBO of Ni and Mo have been performed on JET to systematically analyze the effect of Central RF heating

Main feature:


- ICRH heating: H minority, 5% concentration to heat electrons
- low collisionality ( $\nu_{eff} < 0.2$ )
- about 12 MW NBI
- high central  $q$  to avoid sawteeth
- no total power (NBI + RF) conservation  $\rightarrow$  effect of the RF scan on  $q$ ,  $T_e$ ,  $T_i$ ,  $\omega$  profiles
- no significant MHD activity



The RF power modifies the target plasma affecting mostly Ti , Te , bulk toroidal rotation and q





|       | (MW) |
|-------|------|
| 74354 | 0    |
| 74355 | 1    |
| 74356 | 2    |
| 74359 | 3    |
| 74360 | 3    |
| 74363 | 3    |



ICRH power

Average between  $\rho = 0.2$  and  $\rho = 0.3$






3 MW ICRH


1 MW ICRH

NO ICRH




68383 – marginal H mode  
Low collisionality,  
Similar triangularity  
and elongation


*Average between  $\rho = 0.2$  and  $\rho = 0.3$*




*Average between  $\rho = 0.2$  and  $\rho = 0.3$*

Neoclassical effect?













GS2 (linear, electrostatic) analysis shows trends in the right direction, but never provides an inversion of  $v/D$  (which may occur for ITG  $\rightarrow$  TEM dominant mode transition, obtainable for instance with much lower  $T_i$  gradients than in JET experiments - see e.g. Angioni et al. PPCF 49, 2007);

Tested also the addition of a hot H species (to simulate energetic ion tails after ICRH injection).


The impressive good correlation between  $v/D$  and  $R/LT_i$  suggest a strong neoclassical contribution

$$v_{neo} / D_{neo} = Z_i \cdot (1/n \cdot dn/dr - H \cdot 1/T \cdot dT/dr) ; \quad H \text{ approx } 1$$

Flat density profiles and peaked ion temperature profiles screen impurities out





Simulation of SXR emission with various combinations of transport parameters



- Ni and Mo feature peaked profiles in JET H mode plasmas at low collisionality without sawtooth activity
- Ni and Mo profiles may be made flat-hollow by applying 2-3 MW ICRH
- Impurity pump out has not been explained by turbulence calculations by GS2  
( too large  $R/L_{Ti}$  to get outward turbulent flux)
- Very good correlation between  $v/D$  (Ni) and  $R/L_{Ti}$  , however absolute neoclassical values do not fit the experiment

$\rho = 0.2 - 0.3$




Toroidal rotation


**AUG**

(Courtesy of T. Pütterich, EFPW 2009, Hungary)


**NBI**



**ECRH**




**ICRH**



RF ( electron heating) pump out effect observed in several other experiments:

C-mod, DIII-D, JT-60, TCV and also in non axisymmetric devices (W7-AS)



74357 NO ICRH

74362 3 MW ICRH


ICRH power





For the experimental ne and Te profiles of #68383 at  $r=0.35$ ,  $R/L_{Ti} < 3$  would be necessary for the flow inversion.



CONSORZIO RFX  
Ricerca Formazione Innovazione



L Carraro, C Angoni et al  
EPS 2007 Warsaw



←  $\rho=0.2$  (●)

←  $\rho=0.5$  (►)

- GS2 (linear, electrostatic) scan analysis shows trends in the right direction, but never provides an inversion of  $v/D$  (which it can be for ITG  $\rightarrow$  TEM dominant mode transition reachable, for instance, for much lower Ti gradients - see e.g. Angioni et al. PPCF 49, 2007);


- Tested the addition of a hot H species (to simulate energetic ion tails after ICRH injection). The nature of turbulence remains ITG-dominated: the dominant mode frequency only slightly decreases for increasing beam energies, it never changes sign.


Shots 53548 , 53015 **WITH ICRH**:  
convection may become outward

- Core diffusion decreases
- Core convection also decreases and may become outward

**NO ICRH**

Shot 52136: Strong  
inward convection






- Turbulent transport in center small, dependent on heating
- Central ECH increases diffusion and reduces inward pinch
- Outward convection with ECH real?
- Effect of ICRH similar but less pronounced

Dux et al., PPCF 45, 2003

Courtesy of T. Pütterich @ EFPW 2009, Hungary<sup>27</sup>

TTG-TTF meeting, Cordoba September 2010



Soft-X rays: a vertical camera with  
34 l-o-s (250 $\mu$  filter)

and

a horizontal camera with 17  
channels (350 $\mu$  filter).